

## The Prevalence of Dental Caries among Afghan adults at Dentistry Teaching Clinic, Kabul

Wafiullah Amini<sup>1</sup>, Naseer Ahmad Shinwari<sup>2</sup>, Ahmad Mujtaba Barekzai<sup>3,4,5\*</sup>

<sup>1</sup>Director of Clinic Department of Stomatology Spinghar Institute of Higher Education, Kabul Campus, Afghanistan

<sup>2</sup>Faculty member, Department of Biotechnology and Microbial Technology, Spinghar Institute of Higher Education, Kabul Campus. Afghanistan

<sup>3</sup>Research Director, Spinghar Institute of Higher Education, Kabul Campus, Afghanistan

<sup>4</sup>Faculty Member, Department of Public Health, Spinghar Institute of Higher Education, Kabul Campus, Afghanistan

<sup>5</sup>Technical Manager of Food Safety, providing quality and quantity inspection services for UN-WFP, KIC, Afghanistan

\* **Corresponding Author:** Ahmad Mujtaba Barekzai, Research Director, Faculty member, Department of Public Health, Spinghar University, Kabul Campus, Afghanistan.

**Received date:** 12 June 2022; **Accepted date:** 16 July 2022; **Published date:** 20 July 2022

**Citation:** Amini W, Shinwari NA, Barekzai AM (2022) The Prevalence of Dental Caries among Afghan adults at Dentistry Teaching Clinic, Kabul. Journal of Dental Research and Dental Prospects 01(01): <https://doi.org/10.38207/JDRDP/2022/JUL010105>

**Copyright:** © 2022 Ahmad Mujtaba Barekzai. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

**Objectives:** Caries is a chronic dental disease that causes the loss of hard tooth tissue. If it is not diagnosed and treated in time, it will cause tooth loss anxiety in the act of chewing and speaking affecting the burden and beauty of a person, in addition, it can cause another disease of the teeth and maxillofacial areas.

**Method:** The general purpose of this study is to obtain sufficient information about the prevalence of dental caries for its prevention and treatment so that the occurrence of ocular disease can be controlled and its etiologic should be considered. The study was conducted from the first date of July to the 30 of December among 18-65 years old people referred to the dentistry teaching clinic. In this study, the prevalence of dental caries at different ages was observed in 3440 in 18-65 years old people who were referred to the teaching clinic of dentistry.

**Result:** The prevalence of dental caries in the above ages was 95.98 %, observed in 59.6 % of females and 40.4 % of males with an average of 18-65 years old. The study also found that mood teeth were more caries than old teeth, with a percentage of 85.70. No significant difference was seen in the upper and lower jaws, and the nine age divisions in the group 18-29 years were the highest prevalence of caries with a percentage of 49,08.

**Conclusion:** We can say with great conviction that the prevalence of caries in females compared to males are higher, and it showed that no significant difference was seen in the upper and lower jaws

**Keywords:** chronic dental disease, dental, an observational study

### Introduction

Dental caries is one of the most common chronic dental diseases worldwide. Dental Caries, otherwise known as tooth decay. If it is not diagnosed and treated in time, it will cause tooth loss anxiety in the act of chewing and speaking affecting the burden and beauty of a person [1-3]. In addition, it can cause other diseases of the teeth and maxillofacial areas [4]. In 1997 dental caries were in 190 countries that are part of WHO regions, the average was 2.11 % [5]. According to a cross-sectional survey, approximately 91 % of U.S adults aged 20-64 had dental caries in permanent teeth, dental caries among adults aged 35-64 was 94 % [6]. A Dental Health Survey in the UK shows almost a third of the dentate (31 %) had visible caries in the dentine, representing many millions of people with decay, among the dentate adult with caries at this level, a mean of 2.7 teeth were affected, in particular, 36 % of 25-34 years olds were affected, 26 % of 45-54 years old and 22 % of 65-74 years old [7]. Additionally, a cross-sectional study shows in total, 431 Iranian participants (95.8 %) aged 40-45 years, 163 participants (37.8 %) did not have and decayed

tooth, and 268 participants (62.2 %) were diagnosed with at least one decayed tooth [8]. The case study of 104 Indian shows, that the occurrence of dental caries was found to be slightly higher in females (51.45 %) than males (49.51 %), the occurrence of dental caries was high in vegetarian (85.57 %), non-vegetarians (6.73 %), tobacco user (7.69 %) and the incidence among different age group shown the occurrence of dental caries among 21-30 years' population (26.92 %) [9]. The cross-sectional study in Karachi comprising 500 patients showed that the prevalence of dental caries in Rural 221 patients was (52.7 %), in Urban 198 patients were (47.3 %), and the study reported 3.7 % at the age of 35-34 years [10].

Various factors that have been involved in dental caries are bacteria, genetics, saliva, diet, age, sex, smoking, medicine, and tooth location. According to recent studies, bacteria, colonization by mutant streptococci, and other cariogenic bacteria at a young age could be a key risk factor for caries development. However, the role of mutant streptococci as the main cause of caries has not been proven, because

of the complexity of the oral microflora, which contains several hundred species of bacteria and millions of cells growing on a single tooth surface, no single bacterial species can predict caries development in a particular person. Social-economic status is a stronger predictor of caries risk in children than in adults, it is still important in adults. Because dental caries generally is more prevalent

in lower than higher socioeconomic classes, the dentist should consider the social environment of the patient as available to the patient the medical history, in the analysis of caries risk [11,12]. There is no study demonstrated that Dental caries among Afghan adults. Therefore, we aimed to examine dental caries at the teaching medical hospital in Kabul, Afghanistan.

## Research Methodology

**Study design:** An observational study.

**Study participants:** We selected 3440 patients out of 2034 participants from the random simple selection among 18-65 years old in both gender, which referred to the dentistry teaching clinic (Stomatology of Kabul University of Medical Sciences), from Jul to 30 Dec 2018. Additionally, we examined by examination equipment. The inclusion of our study is both genders, with adults ages 18-65 years, and all patients were allowed to join in our study. And excluded from our study are the patients with the chronic disease under the age

of 18 years old and over 65 years old. As well, we trained the data collector for all variables. Such as sex (phenotypes) we detect the sex of participants by phenotypes. The age we detect the age of participants between 18-65 years old. We detect the Maxilla's with anatomic location of participants and detect the Salfi jaws with anatomic location of participants. We detect the anterior and posterior teeth by anatomic location of participants. The study was ethically approved by the medical bioethics committee of Spinghar Institute of Higher Education (Code: 1386-1405)

## Results

The prevalence of dental caries in both females and males was 3302 tons (95.98%). Of the 3440 patients, 1402 were males and 2038 were females. In the male class, out of 1402 of them, 1332 were caries. Of the 2038 female patients, the caries outbreak was reported in 1970. In this study which was observed in 95.98 % of caries patients, the prevalence of dental caries was different in different individuals for some clients, caries had more than one tooth. Among the teethinfected with caries, 4349 teeth infected with caries were observed and recorded from both anterior and posterior teeth of the upper and lower jaws. 623 (14.30 %) anterior teeth and 3735 (85.70 %) posterior

teeth were exposed to caries. Caries outbreaks were recorded in the upper jaw 2216 (50.85 %) and the lower jaw 2142 teeth (49.15 %). The study of the prevalence of caries age in terms of age was conducted in five age groups. The first group, which was between the ages of 18-29, 2139 teeth 49.08 %, the second group, which was between the ages of 30-39, 889 teeth 20.39 %, the third group, which was between the ages of 40-49, 740 teeth 16.98 %, the fourth group, which was between the ages of 50-59, 379 teeth 8.69 %, the fifth group, which was 60 years old and above 211 teeth which were 4.84 % registered.

**Table 1:** The demographics of participants:

Variables		Percentage	Frequency
Sex	Female	59.6%	1970
	Male	40.4%	1332
Caries Location on jaw	Maxilla	50.85	2216
	Mandibula	49.15	2142
Caries Location	Anterior teeth	14.30	623
	Posterior teeth	85.70	3735
Age	18-29 years old	49.08	2139
	30-39 years old	20.39	889
	40-49 years old	16.98	740
	50-59 years old	8.69	379
	60 years old and older	4.84	211

## Discussion

Our study demonstrated that the prevalence of dental caries in both females and males was 3302 tons (95.98 %). Of the 3440 patients, 1402 were males and 2038 were females. In the male class, out of 1402 of them, 1332 were caries. Of the 2038 female patients, the caries outbreak was reported in 1970. In this study which was

observed in 95.98 % of caries patients, the prevalence of dental caries was different in different individuals for some clients, caries had more than one tooth. These findings support our aim. For instance, in a study conducted by Devey Bruce et al in the United States in 2011-2012, the prevalence of caries was 91 % which is almost consistent

with our results [13]. Also, in our study, the highest incidence of caries was found in females at 59.6 %, According to research literature conducted by social Karpal Singh and colleagues in Tanzania in 2013, more than 436 patients received class at 58.7 %, which is more than the male class that complies [14]. In our research, there was no significant difference between Alavi and Salafi jaws (49.15 % of Salafi jaws and 50.85 % of Maxillas). According to the research literature conducted by Sohal Karpla Singh and colleagues in Tanzania in 2013, no difference was found in terms of the jaw (50.3 % of the Maxilla and 49.7 % of the Salafi jaw), which is in perfect agreement. The research I did on the posterior teeth found that the highest incidence of caries was 85.70 % and that of the anterior teeth was 14.30 %, as seen in the literature, in a study conducted by Dr. Mohammad Usman Riyadh, they found that 77 % of Caries were found in the posterior teeth and 23 % in the anterior teeth, it can be seen that in both studies, posterior teeth show the highest incidence

### Conclusion

We can say with great conviction that the prevalence of caries in females compared to males are higher, and it showed that no significant difference was seen in the upper and lower jaws

**Acknowledgments:** We all thank full of the data collection team and participants.

**Data availability statement:** The raw data supporting the conclusions of this article will be made available by the authors, on reasonable request to the corresponding author.

**Competing interests:** All authors declared no potential personal or financial conflicts of interest.

### References

- Selwitz RH, Ismail AI, Pitts NB (2007) Dental caries. *The Lancet*. 369(9555): 51-9.
- Veiga NJ, Aires D, Douglas F, Pereira M, Vaz A, et al. (2016) Dental caries: A review. *Journal of Dental and Oral Health*. 2(5): 1-3.
- Bagramian RA, Garcia-Godoy F, Volpe AR (2009) The global increase in dental caries. A pending public health crisis. *Am J dent*. 22(1): 3-8.
- Pitts NB, Zero DT, Marsh PD, Ekstrand K, Weintraub JA, et al. (2017) Dental caries. *Nature reviews Disease primers*. 3(1): 1-16.
- da Silveira Moreira R (2012) Epidemiology of dental caries in the world. *Oral Health Care.-Pediatric, Research, Epidemiology and Clinical Practices*.
- Dye B, Thornton-Evans G, Li X, Iafolla T (2015) Dental caries and tooth loss in adults in the United States, 2011-2012. *NCHS Data Brief*. (197): 197.
- White DA, Tsakos G, Pitts NB, Fuller E, Douglas GV, et al. (2012) Adult Dental Health Survey 2009: common oral health

of caries. From 2001-to 2004 in Istanbul, Turkey, also the prevalence of caries in posterior teeth compared to anterior teeth had been confirmed, with anterior teeth at 31.6 % and posterior teeth at 68.4 %. According to the age group of caries in the group 18-29 years, the highest incidence of caries in our study was 49.08 %. In a study conducted in Tanzania by Sohal Karpal Singh et al, the incidence of caries in the age group of 18-29 years received 41.1 %, which is the highest incidence of caries among other groups, in our research, caries in this group also lot. The result of this study shows that the prevalence of caries is higher in females than in males and in the location of the teeth in the jaws of posterior teeth compared to anterior teeth. And no significant difference was seen from the jawbone, its events still show the most cases according to the group of 18-29 years. The prevalence of dental caries in our center (22.6 %) falls within the previously reported 4-30 % rates for Nigeria.

**Ethics statement:** This study was ethically approved by the medical bioethics committee of the SIHE ethics committee (code: 1386-1405). The patients/participants provided their written informed consent to participate in this study.

**Author contributions:** WA, NAS was involved in the study's conception, design, statistical analysis, and interpretation of the data. AMB, and NAS, were involved in data collection, data cleaning, statistical analysis, and manuscript drafting. AMB supervised the study. All authors approved the final manuscript for submission.

**Funding:** N/A

conditions and their impact on the population. *British dental journal*. 213(11): 567-72.

- Conrad J, Retelsdorf J, Attia S, Dörfer C, Mekhemar M (2020) German dentists' preferences for the treatment of apical periodontitis: a cross-sectional survey. *International Journal of Environmental Research and Public Health*. 17(20): 7447.
- Khan AA, Jain SK, Shrivastav A (2008) Prevalence of dental caries among the population of Gwalior (India) in relation of different associated factors. *European journal of dentistry*. 2(02): 081-5.
- Siddiqui TM (2013) An Epidemiological Study of Prevalence of Dental Caries and Periodontal Disease among Adults in Deprived Areas-Karachi. *JOHR*.
- Bowden G (2000) The microbial ecology of dental caries. *Microbial ecology in health and disease*. 12(3): 138-48.
- Kutsch VK (2014) Dental caries: an updated medical model of risk assessment. *The Journal of prosthetic dentistry*. 111(4): 280-5.

13. Ross CT (2015) A multi-level Bayesian analysis of racial bias in police shootings at the county-level in the United States, 2011–2014. PloS one. 10(11): e0141854.
14. Sohal KS, Kalyanyama BM, Simon EN (2020) The pattern of assault-related oral and maxillofacial injuries among patients treated at the Muhimbili National Hospital, Dar es Salaam, Tanzania. t Med. 2(2): 119-124.