

The Continuing Trend Of Sodium Nitrite Suicides: A Case Series And Brief Literature Review To Guide Diagnosis

Dr Leilah Bidwell, MBBS, MCLinEd*

Princess Alexandra Hospital, Harlow, UK

***Corresponding Author:** Dr Leilah Bidwell, Princess Alexandra Hospital, Harlow, UK

Received date: 02 November 2025; **Accepted date:** 11 November 2025; **Published date:** 13 November 2025

Citation: Bidwell L (2025) The Continuing Trend Of Sodium Nitrite Suicides: A Case Series And Brief Literature Review To Guide Diagnosis. J Med Case Rep Case Series 6(06): <https://doi.org/10.38207/JMCRCS/2025/NOV06060566>

Copyright: © 2025 Leilah Bidwell. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Introduction: Sodium nitrite (NaNO_2) has worryingly emerged as an increasingly used method of suicide due to its ready availability and misleading online portrayal as a peaceful means of death. Recent media coverage, including a Channel 4 documentary, has drawn attention to a growing number of cases.

Methods: We report three sodium-nitrite-related deaths occurring over a nine-month period (May 2024 – February 2025) in a district hospital in southeast England. Each case was investigated post-mortem, with additional toxicological testing for nitrites, nitrates, and methaemoglobin where indicated.

Results: Not all cases displayed classic features such as chocolate-brown blood or organ discolouration. In each, different investigative triggers led to nitrite testing, confirming markedly elevated nitrite levels (up to 30,000-fold above normal). Methemoglobinemia was identified in all cases.

Conclusion: Detection of sodium nitrite and methemoglobinemia post-mortem is challenging due to sample degradation. Early collection and analysis are essential for reliable results. Vigilance among clinicians and pathologists is vital, as no single diagnostic clue is universally present. Ongoing awareness and early toxicological testing are essential to ensure accurate case recognition and monitoring of this evolving trend.

Keywords: Suicide, Sodium Nitrite, NaNO_2 , Case Series, Overdose, Autopsy, Toxicology, Post-mortem

Introduction

Sodium nitrite (NaNO_2) is a white-to-yellow crystalline powder [1] used industrially as an antimicrobial, food preservative, and component of anti-freeze [2, 3]. At therapeutic doses it acts as an antidote to cyanide poisoning [1] but, ingestion of larger quantities causes cellular hypoxia, potent vasodilation and severe hypotension [4]. This is due to acute methoglobinemia where iron oxidation from Fe^{2+} to Fe^{3+} renders MetHb unable to bind to, and therefore transport oxygen to the tissues, usually MetHb accounts for 1-3% of total circulating haemoglobin, once levels rise above 50% patients become symptomatic, and fatality is usually seen at concentrations above 70% [5].

The antidote for nitrite toxicity is methylene blue [4]. Sodium nitrite has an estimated lethal dose of 2.6g, although a fatal case after 1g, and survival after ingestion of 6g have been reported. It is readily available as pure granules via online retailers and use in suicide has been discussed in online forums [6]. Commercial “suicide kits” including antiemetics, antacids together with detailed instructions remain readily accessible online [5, 7, 8, 9].

Toxicology

Testing for nitrites or nitrates is not part of routine toxicology screening postmortem [7, 10]. Commonly documented autopsy

findings include those consistent with methoglobinemia; blue-grey hypostasis and dark brown discolouration of blood and internal organs [1] some studies and case reports have also noted presence of powder residue in the gastrointestinal tract [10].

Regulatory Developments

In June 2021 [4, 5], Italian authorities blocked access to a pro-suicide website to limit online promotion of sodium nitrite use. Governmental bodies have regulated the use of the NaNO_2 to protect public health and safety [10]. The UK Home Office subsequently classified sodium nitrite a reportable substance requiring suspicious purchases to be reported [11]. However, this does not prevent the substance being readily available with some websites offering consultations [4] with guidance on dosage and how to prevent vomiting. Reported sellers known as “The Ukraine Supplier [12]” and other sources recommend sodium nitrite as a peaceful option, such as The Peaceful Pill Handbook [1, 11].

Epidemiology

Multiple case series and reviews report an increasing global trend in sodium nitrite use for suicide [1, 4, 5, 11]. It has been suggested the

most affected demographic are under the age of 30, partially due to the use of the internet, and primarily suicide forums recommending sodium nitrite as a method of suicide [7]. It is also of note that those using this method are in the 'post-covid era' which may have contributed to isolation, and social media influence.

Awareness of sodium nitrite suicides has increased following widely reported cases involving alleged such as Kenneth Law in Canada and Leonid Zakutenko in Ukraine; both have been the subject of ongoing official investigations and media coverage regarding the sale of NaNO_2 to vulnerable individuals who later committed suicide [5]. Mr Law was arrested by Canadian police in May 2022 and is awaiting trial set for January 2026 for 14 counts of first-degree murder [13]. Mr Zakutenko, however, remains in Ukraine, investigations allege he has sold NaNO_2 over 1000 times and caused at least 93 UK deaths [12].

We report three locally identified cases over nine months in a district hospital in southeast England, each discovered through different investigative triggers.

Case Series

Case 1 — May 2024

A 24-year-old African Caribbean male was found unresponsive in a hotel room. There were no external injuries. Two large bags (302 g) of pale-yellow crystalline powder, digital weighing scales, and orange watery vomit were found at the scene.

History: Known anxiety and hypertension; regular fluoxetine 20 mg and prazosin 20 mg.

Findings: External examination revealed white foamy residue on lips and clothing. Internal examination showed left ventricular hypertrophy (24 mm) and pulmonary oedema.

Toxicology: Metoclopramide was detected. Subsequent nitrite analysis revealed a ~30 000-fold elevation in nitrite (NO_2^-) with no detectable nitrate (NO_3^-).

Cause Of Death: Nitrite toxicity causing methemoglobinemia and profound hypotension.

Case 2 — November 2024

A 27-year-old female was found unresponsive; attempted resuscitation had been performed.

Findings: Endotracheal tube in situ, cyanosis of lips and face, and multiple healed self-harm scars. Post-mortem CT showed severe pulmonary oedema and congestive cardiac failure.

Toxicology: No common drugs detected on initial screening. Extended testing showed a 254-fold increase in nitrate levels and elevated methemoglobin.

Cause Of Death: Multi-organ failure secondary to methemoglobinemia from nitrite toxicity.

Case 3 — February 2025

A 25-year-old Caucasian female with autism and a history of self-harm and previous suicide attempts was found collapsed after vomiting was heard during a phone call.

Scene: A handwritten note and packaging labelled 'Sodium Nitrite BASF Food Preservative NaNO_2 (purity 99.6%)' were discovered by relatives.

Findings: Slim build, cyanosis, and scarring consistent with prior self-harm. No internal organ discolouration.

Toxicology: Sertraline, lamotrigine, and naloxone were present at non-toxic concentrations. Nitrite analysis of femoral blood revealed a ~21 000-fold nitrite elevation with no nitrate detected.

Cause Of Death: Nitrite toxicity with methemoglobinemia and hypotension.




Figure 1: photograph

Summary Of Analytical Findings

Parameter	Case 1	Case 2	Case 3
Age/Sex	24 / M	27 / F	25 / F
Trigger for testing	High metoclopramide; unidentified powder	Facial cyanosis; pulmonary oedema	NaNO ₂ found at scene
Time between death and sample processing	8 weeks	8 weeks	8 weeks
Sample type	Unpreserved femoral vein blood sample Stored at 2-4°C	Plain blood sample, undisclosed source Stored at 2-4°C	Unpreserved femoral vein blood sample Stored at 2-4°C
Sample quality	Evidence of red blood cell lysis; amounts shown likely to be substantial underestimate of true concentrations		
Analytical method	Ozone chemiluminescence		
Nitrite (NO ₂ ⁻)	7314µM	5171µM	Not detected
Nitrate (NO ₃ ⁻)	0µM	0µM	7605 µM
MetHb (%)	>70	65	>70
Cause of death	1a. Methemoglobinemia and hypotension 1b. Nitrite toxicity	1a. Multi-organ failure with methaemoglobinaemia 1b. Nitrite toxicity	1a. Methaemoglobinaemia and hypotension 1b. Nitrite toxicity

Analytical Methodology

No stabilisers were used prior to nitrite / nitrate testing. Blood samples from all three cases were centrifuged at 15,000 g, 4°C, for 5 minutes to remove debris, yielding a clear supernatant for dilution. Samples were analysed using gas-phase chemiluminescence (NOA 280i, Sievers) measuring the reaction between Nitric Oxide (NO) and ozone (O₃). For every mole of NO₂⁻ or NO₃⁻ one mole of NO is generated.

Interpretation Of Nitrite Vs Nitrate Findings

In healthy fasting individuals, plasma NO₂⁻ is between 0.1-0.4µM, and for NO₃⁻ is it 20-40µM¹⁴. Dietary NO₃⁻ usually contributes significantly to circulating NO₃⁻ and NO₂⁻ concentrations. Dietary nitrate intake can increase NO₃⁻ ~10-fold and as a consequence of transport through the enterosalivary circuit, a doubling of circulating NO₂⁻ concentrations. Therefore, if there is no measurable NO₃⁻ a dietary source is unlikely to be responsible for the raise in NO₂⁻. NO₂⁻ has a very short plasma half-life (180 seconds) [7] before it is converted to NO₃⁻. This conversion can and does continue postmortem [15]. As such, in the absence of NO₂⁻, NO₃⁻ can be used as an indirect indicator of nitrite intake in suspicious circumstances, although the diagnosis is still based on methaemoglobin analysis [16]. This is because the principal mechanism of nitrite toxicity is the oxidation of the ferrous iron Fe²⁺ to Fe³⁺ producing methaemoglobin [17]. Under normal conditions approximately 3% of haemoglobin is oxidised to methaemoglobin [15], in cases of nitrite ingestion this is substantially higher.

Discussion

Clinical Presentation

These three cases illustrate the variable clinical and postmortem presentation of sodium nitrite toxicity. Characteristic signs such as darkened blood may be absent. Alternate indicators: the presence of anti-emetics, powder residues, or unexplained cyanosis were key to initiating further toxicological testing.

In these deaths recognition depended on indirect indicators: antiemetic presence, unexplained cyanosis, or scene evidence of NaNO₂.

Analytical And Diagnostic Considerations

Analytical confirmation remains challenging, as nitrite concentrations degrade rapidly postmortem, with conversion to nitrate continuing after death [4]. Studies indicate up to 95% of the original concentration can be lost within one hour at room temperature, emphasising the importance of prompt collection. However, nitrate levels can be measured to compensate for this [7]. Nitrite stability can be preserved at -20°C if stabilised with potassium ferricyanide [11], however, this requires an initial suspicion as this is not a routine storage or preservative method. Interestingly one study found urinalysis often tests positive for nitrites without evidence of urinary tract infection in patients [1] in those using NaNO₂ as a suicide method, however this was only used in the last 8 cases. This was more reliable in females, with an assumed sensitivity of 45% and specificity of 85% overall. Ultimately nitrate, nitrite and methaemoglobin levels should be tested immediately to prevent misinterpretation of values [11].

Epidemiological And Public-Health Implications

The suggestion that this is becoming a new trend is compounded by a 30% increase reported in suspected self-poisoning suicide attempts among individuals aged 10-19 in United States as reported by the National Poison Data System (NPDS) [19]. Media coverage and online availability likely contribute to its growing use [1].

The identification of three confirmed cases within nine months at a single district hospital raises concern for potential under-recognition elsewhere. However, this is a local observation and cannot be interpreted as evidence of national prevalence.

Preventive And Regulatory Actions

Regulatory responses are emerging. The UK Home Office classification of NaNO₂ as a 'reportable substance [10]' and the Italian judicial blocking of pro-suicide sites are positive steps. [4, 5]. Nonetheless, the compound remains accessible through international suppliers [1, 11] and online forums, underscoring the need for coordinated global regulation and public health surveillance to limit further harm.

Forensic Diagnostic Implications

While studies and previous case reports have documented common autopsy findings, these are not always present. The most recent systematic review found 19/63 cases had dark discolouration of the blood [5].

A case series of 20 patients found 80% had a history of depression or mental health issues [9], and anti-emetic drugs were present in 35% of cases. Other studies found evidence of NaNO₂ at the scene in most cases [1, 10], either packaging, a glass, or white powder residue.

It is important to test as early as possible if any suspicion of Nitrite involved deaths. Detection of packaging or powder residue at the scene remains one of the most consistent clues [1, 10]. Femoral blood is considered the most reliable sample for nitrite analysis [5]. Prompt sampling, along with parallel measurement of nitrates and methaemoglobin, provides the best diagnostic accuracy. Studies monitoring methaemoglobin changes in samples during storage recommend samples be taken at autopsy [18], refrigerated and stored in EDTA preservative, and for longer storage samples should be frozen with cryoprotectant and stored at -80°C.

References

- Stephenson L, Wills S, van den Heuvel C, Humphries M, Byard RW (2022) Increasing use of sodium nitrite in suicides-an emerging trend. *Forensic Sci Med Pathol*. 18(3): 311-318.
- Olson KR, Benowitz NL, Anderson IB, Blanc PD, Clark RF, et al. (2018) Poisoning and drug overdose. 7. New York: McGraw Hill Education (LANGE). 339-340.
- Laue W, Thiemann M, Scheibler E, Wiegand KW (2000) Nitrates and nitrites. *Ullmann's Encyclopedia of Industrial Chemistry*.

Limitations

This series is limited by its small sample size, single institution setting, and potential ascertainment bias, as cases of sodium nitrite toxicity may be underreported or misclassified. Additionally, varying postmortem intervals and testing conditions may have influenced nitrite stability and measured concentrations. The findings are descriptive and cannot be generalised beyond the local population.

Conclusion

Sodium nitrite toxicity represents a growing but under-recognised method of self-poisoning. The variability in clinical and post-mortem findings underscores the importance of considering NaNO₂ ingestion, even if classic features are absent. Early sampling, preservation, and methaemoglobin testing are essential for reliable post-mortem diagnosis.

This local three-case series demonstrates diagnostic and analytical challenges and suggests possible under-recognition of similar deaths. Interpretation should be cautious due to small sample size, single-centre data, and potential bias. Broader multicentre surveillance and regulatory collaboration are needed to accurately characterise this emerging trend and inform prevention strategies.

Author Contributions: With thanks to contributions from Dr Angela Cymerman, Dr Vasi Sundaresan and Dr Preethi Gopinath for access to PM case reports.

Ethical Declarations

Funding: This research received no external funding.

Informed Consent Statement: No informed consent was required, as patients are not identifiable from the information presented in this case series.

Ethical Compliance: All tissue and samples were handled according to the Human Tissue Act (2004) during and after post-mortem examination.

Conflicts of Interest: The author declares no conflicts of interest.

6. Wettstein ZS, Yarid NA, Shah S (2022) Fatal methaemoglobinemia due to intentional sodium nitrite ingestion. *BMJ Case Reports*. 15(12): e252954.
7. Caparrelli V, Pallocci M, Tittarelli R, Russo C, Donato L, et al. (2025) Sodium Nitrite-Related Fatalities: Are We Facing a New Trend? Case Report and Literature Review. *Forensic Sciences*. 5(3): 42.
8. Durão C, Pedrosa F, Dinis-Oliveira RJ (2020) A fatal case by a suicide kit containing sodium nitrite ordered on the internet. *J. Forensic Leg. Med.* 73: 101989.
9. Hikin LJ, Ho J, Morley SR, Ahluwalia A, Smith PR (2023) Sodium nitrite poisoning: A series of 20 fatalities in which post-mortem blood nitrite and nitrate concentrations are reported. *Forensic Science International*. 345: 111610.
10. Padovano M, Aromatario M, D'Errico S, Concato M, Manetti F, et al. (2022) Sodium Nitrite Intoxication and Death: Summarizing Evidence to Facilitate Diagnosis. *International Journal of Environmental Research and Public Health*. 19(21): 13996.
11. Kaubrytė SS, Chmieliauskas S, Salyklytė G, Laima S, Vasiljevaitė D, et al. (2025) Fatal Outcome of Suicidal Multi-Substance Ingestion Involving Sodium Nitrate and Nitrite Toxicity: A Case Report and Literature Review. *Acta medica Lituanica*. 32(1): 160–172.
12. Smith T, Crawford A (2024) How we tracked down the Ukrainian poison seller. *BBC News*.
13. Yousif N (2024) Canadian 'poison killer' Kenneth Law to head straight to trial. *BBC News*.
14. Bescos R, Gallardo-Alfaro L, Ashor A, Rizzolo-Brime L, Siervo M et al. (2024) Nitrate and nitrite bioavailability in plasma and saliva: their association with blood pressure-A systematic review and meta-analysis. *Free Radical Biology and Medicine*. 226: 70-83.
15. Kim M, Kim S, Yang W, Sim J (2022) Determination of Nitrite and Nitrate in Postmortem Whole Blood Samples of 10 Sodium Nitrite Poisoning Cases: The Importance of Nitrate in Determining Nitrite Poisoning. *Forensic Science International*. 335: 111279.
16. Taus F, Pigaiani N, Bortolotti F, Mazzoleni G, Brevi M, et al. (2021) Direct and specific analysis of nitrite and nitrate in biological and non-biological samples by capillary ion analysis for the rapid identification of fatal intoxications with sodium nitrite. *Forensic Science International*. 325: 110855.
17. Institute of Medicine (1995) *Environmental Medicine: Integrating a Missing Element into Medical Education*. National Academies.
18. Varlet V, Ryser E, Augsburg M, Palmiere C (2018) Stability of postmortem methemoglobin: Artifactual changes caused by storage conditions. *Forensic Science International*. 283: 21–28.
19. Farah R (2023) Suspected Suicide Attempts by Self-Poisoning Among Persons Aged 10–19 Years During the COVID-19 Pandemic-United States, 2020–2022. *Morbidity and Mortality Weekly Report (MMWR)*. 72(16): 426-430.